

American Society of Anaesthesiologist (ASA) classification


Class

- 1 Healthy patient, no medical problems
- 2 Mild systemic disease eg hypertension
- 3 Severe systemic disease, but is not incapacitating
- 4 Severe systemic disease that is a constant threat of life
- 5 Moribund expected to live < 24 hours irrespective of operation
- E Operation is an emergency

Predictors of difficult intubation

Mallampatti Score III & IV
 Inability to open mouth > 4 cm
 Thyro-mental distances < 6 cm
 Limitation of neck movement
 Difficulty of protruding lower jaw
 History of difficult intubation

Mallampati Score



Sedation Score

6	Inadequate	Anxious, agitated, or in pain
5	Minimal	Spontaneously awake without stimulus
4	Moderate	Drowsy, eyes open or closed, easily roused by verbal stimulus
3	Moderate / deep	Rouses with moderate tactile, loud verbal stimulus
2	Deep	Rouses slowly to consciousness with painful stimulus
1	Deep	Rouses, but not to consciousness, with painful stimulus
0	Anaesthesia	Unresponsive to painful stimulus

Minimum home discharge criteria guidelines

Medical observations satisfactory
 Resumption of pre-sedation level of function
 Ability to walk at pre-sedation level safely
 Able to tolerate oral intake
 Discharge information given