



# Patient Oxygen Delivery and Corresponding PPE

## DROPLET PRECAUTIONS

### Oxygen Delivery System And Flow Rate (L/min)



Nasal Prongs: 1-4L



Hudson Mask: 4-8L



Non-rebreather Mask: 8-15L

**HFNP are currently contraindicated in confirmed and suspected COVID 19 patients outside of the ICU setting**



High Flow Nasal Prongs

### Patient PPE



Surgical Mask  
-worn at all times including **transfers**



Full Face Shield  
- worn only during prolonged procedures or **transfers**



Full Face Shield  
- worn only when staff are in room or during **transfers**

### Staff PPE



Full length gown, surgical mask, hat, glasses, sterile gloves

## AEROSOL PRECAUTIONS

**\*\*\*HIGH RISK\*\*\***

**ONLY TO BE PERFORMED IN THE AEROSOL GENERATING PROCEDURE ROOMS.  
CONSIDER IF NECESSARY MANAGEMENT PRIOR TO WARD TRANSFER.**

**Nebulisers and NIV are contraindicated in confirmed and suspected COVID 19 patients  
For non-COVID 19 patients requiring NIV/neb, use the following:**



Nebuliser



NIV



Intubation



Full Face Shield  
-worn during **transfer**

Not Applicable



Full length gown, surgical hood, N95 mask, full face shield, sterile gloves